arXiv:1612.04543v1 [math.DG] 14 Dec 2016

Locally conformally cocalibrated G_2-structures

Arezoo Zohrabi

Dipartamento di matematica, Università degli studenti di Torino, Italy

Abstract

We study the condition in which G_2-structures are introduced by a non closed four-form, although they are satisfying locally conformal conditions.All solutions are found in the case when the Lee form of G_2-structures is non-zero and $\mathfrak{g}$ introduces seven-dimensional Lie algebras, The main results are given in preposition1 and theorem1.

Contents

1 Nearly Half Flat SU(3)-structures **3**

2 Locally conformal cocalibrated G_2-structures **4**

3 Associated Lie algebras **7**

Introduction

A G_2-structure (M^7, φ) on a seven-dimensional manifold M^7 can be characterized by the existence of globally 3-form φ called fundamental 3-form. the classes of G_2-structures can be described in the terms of the exterior derivatives of fundamental 3-form φ and the 4-form $\phi = *\varphi$, where $*$ is the Hodge operator defined from metric and the derivative of V_{g_φ}, in this paper we focus our attention on the class of locally conformally cocalibrated G_2-structures (usually denoted by L.C.CC G_2-structures), which are characterized by the condition $d\phi = -\theta \wedge \phi$ for a closed non vanishing 1-form θ also known as the Lee form of the G_2-structures. Let M be connected n-dimensional manifold with a Riemannian metric on M, Let $\bigtriangledown$ be the Levi-Civita connection of G and for $p, q \in M$ joint by smooth path γ then parallel transport along γ using $\bigtriangledown$ define an isometry between the tangent space of T_pM , T_qM.
A seven-dimensional connected,oriented Riemannian manifold M^7 with the Holonomy [1]contained in G_2 is characterized by the existence of a G_2 reduction of its orthogonal frame bundle with the restriction of Cartan form of the Levi-Civita connection, alternatively can be seen that $G_2 \subset SO(7)$ is defined as the stabilizer under the action of $GL(7, \mathbb{R})$ of stable 3-form $\varphi\,'$ or equivalently 4-form ϕ' which are related $\phi' = *\varphi\,'$
A seven-dimensional nilmanifold is a compact manifold which corresponding lie group G acts transitively on and in homogeneous case G/Γ where Γ is a discrete subgroup whit the property for $a.g \in G/\Gamma \Longrightarrow b(a.g) = (ba).g \in G/\Gamma$ for $a, b \in \Gamma$ and $g \in G$, equivalently Lie group G can define a nilmanifold if and only if its lie algebra $\mathfrak{g}$ admits a basis $\{e_1, e_2, ..., e_7\}$ and invariant forms on a homogeneous case are determined by forms on $\mathfrak{g}$, so we can consider seven-dimensional nilpotent Lie algebras consequently.
G_2 is a compact lie group as a closed subgroup of the orthogonal group $O(7, \mathbb{R}) = \{Q \in GL(n, \mathbb{R}) | Q^T Q = QQ^T = I\}$, $G_2 = \{g \in GL(n, \mathbb{R}) | g^*(\phi) = \phi\}$ a few properties of G_2 will be used here. The group G_2 acts irreducibly on $\mathbb{R}^7$ and acts transitively on the unit sphere $S^6 \subset \mathbb{R}^7$ and preserve the metric and orientation for which the basis $\{e_1, e_2, ..., e_7\}$ is an oriented orthonormal basis, the notation g and $\langle ., . \rangle$ will be used to refer the metric.
The stabilizer subgroup of any non-zero vector in $\mathbb{R}^7$ is isomorphism to $SU(3) \subset SO(6)$, so that $S^6 = G_2/SU(3)$. Since SU(3) acts transitively on $S^5 \subset \mathbb{R}^6$, it follows that G_2 acts transitively on the set of orthonormal pairs of vectors in $\mathbb{R}^7$. However, G_2 does not act transitively on the set of orthonormal triples of vectors in $\mathbb{R}^7$ since it preserve the 3-form $\varphi = *\phi$.
By definitions, G_2 is a compact lie group as a closed subgroup of the orthog-

[1]The holonomy group of metric g is the group of isometries of T_pM generated by parallel transport around closed loop at the point p, i.e. $Hol(g) \leq O(n)$

onal group $O(7,\mathbb{R})$ and Lie algebra of G_2 is $\mathfrak{g}_2$.

1 Nearly Half Flat SU(3)-structures

Definition. Let $\mathfrak{g}$ a lie algebra then $\mathfrak{g}$ is nilpotent if the lower central series terminates , i.e if $\mathfrak{g}_n = 0$ for some $n \in \mathbb{N}$. this means that

$$[X_1, X_2, [...[X_n, Y]...]] = ad_{X_1} ad_{X_2} ... ad_{X_n} Y \in \mathfrak{g}_n$$

$$X_1, X_2, ..., X_n, Y \in \mathfrak{g}$$

so that $ad_{X_1} ad_{X_2} ... ad_{X_n} = 0$. Note that if $\mathfrak{gl}(n,\mathbb{K})$ is the set of $n \times n$ with the entries of $\mathbb{K}$, then the subgroup consisting of strictly upper triangular matrices is nilpotent lie algebras.
A Lie derivative for a general differential form is likewise a contraction, taking into account the variant in X : $L_x w = \iota_x dw + d(\iota_x w)$ knowing as Cartan's Formula.

Definition. For a lie algebra $\mathfrak{h}$ of dimension 2m, a closed two-form $\omega \in \Lambda^2 \mathfrak{h}^*$ is called symplectic if it is non-degenerate i.e. in the case of $m = 3$ $\omega^3 \neq 0$.

Note1. From the sixtieth classes of G_2-structures of in this paper we consider the class of torsion $W_1 \oplus W_3 \oplus W_4$.
Gray and Harvella in [4] proved that there exist sixteen different class of almost Hermitian structures attending to the behavior of the covariant derivative of its *Käler* form. equivalently the different classes of SU(3)-structures can be defined in terms of form ω, ψ_+ and ψ_- in particular we are interested on Nearly Half-Flat which is defined on six-dimension lie algebra $\mathfrak{h}$.
An SU(3)-structure on a manifold of real dimension 6 consists of a Hermitian structure (g, J, ω) and a $(0,3)-$form ψ; and SU(3) is the stabilizer of the transitive action of the group G_2 on S_6.

Definition. We call an SU(3)-structures (ω, ψ_+, ψ_-) is nearly half-flat on a 6-dimensional manifold when it satisfied the equation:

$$d\psi_- = \frac{1}{2}(\omega \wedge \omega)$$

However six-dimensional nilpotent lie algebras admitting a nearly half-flat SU(3)-structures are not classified,Those which admitting a double half-flat are classified by Choissi and Swann in [1] and as double half-flat are in particular nearly half-flat this allows one to shows existence locally conformally cocalibrated G_2-structure.

2 Locally conformal cocalibrated G_2-structures

Now we consider the L.C.CC G_2-structure which is introduce by a 4-form ϕ s.t. $d\phi = \theta \wedge \phi$ for a closed 1-form θ, and $\phi \in \Lambda^4 V^*$ compatible with orientation of $\mathfrak{g}$ and the underlying metric g, consider the $X \in \mathfrak{z}(\mathfrak{h})$, $L_X = 0$ where $[X, Y] = L_X Y$ $X = \theta^\sharp$ is isomorphism induced by the metric between the space and dual space, where:

$$d\phi = \theta \wedge \phi, X = \theta^\sharp \in \mathfrak{g}$$

s.t. $\phi = \sigma + \psi_- \wedge \theta$, $\iota_X \omega^2 = 0$ and $\iota_Y \psi_- = 0 \Rightarrow \phi = \theta \wedge \phi$
$\Rightarrow d\omega^2 = (\omega^2 - d\psi_-) \wedge \alpha$

observe that $\theta(X/|X|) = 1$ for $X \in \mathfrak{z}(\mathfrak{h})$ and $L_X = 0$, $[X, Y] = L_X Y$

where $|X| = \sqrt{g(X,Y)}$ for $X^\perp = \{Y \in \mathfrak{g} \mid g(X,Y) = 0\}$,

$$\theta = \theta_1 e^1 + \theta_2 e^2 + ... + \theta_6 e^6 + \theta_7 e^7$$

$L_X \phi = \iota_X d\phi + d\iota_X \phi$, $L_X \phi = \phi + d(-\psi_-)$ where $\psi_- = -\iota_X \phi$ where pair (ω, ψ_-) is defining a neraly half-falat SU(3)-structure,
$de^i(e_l, e_k) = -e^i([e_l, e_k])$, $X = \theta^\sharp$, $\theta(Y) = g(X,Y) = \theta^\sharp for \forall Y \in \mathfrak{g}$ theta is dual form of vector, s.t. $\theta \in \Lambda^1 \mathfrak{g}^*$ our vector $X \in [\mathfrak{g}, \mathfrak{g}]$ where $\Lambda^1 \mathfrak{g} \longrightarrow \Lambda^1 \mathfrak{g}^*$, $X \in [\mathfrak{g}, \mathfrak{g}]$ and G_2 acts irreducibly on V and hence on $\Lambda^1 V^*$ where there is a short exact sequence:

$$0 \longrightarrow \mathbb{R}X \longrightarrow \mathfrak{g} \longrightarrow \mathfrak{h} \longrightarrow 0$$

with the definition of this homomorphism $\pi^* : \Lambda^k \mathfrak{g}^* \longrightarrow \Lambda^1 \mathfrak{h}^*$ it send $\pi^* \alpha$ for one-form α s.t.

$$\pi^*[X, Y] = [\pi^* X, \pi^* Y]$$

Proposition 1 . *Consider ϕ be a 4-form Locally Conformal Cocalibrated G_2-structures i.e. $d\phi = \theta \wedge \phi$,$d\theta = 0$, $\theta \in \Lambda^1\mathfrak{g}^*$ and the pair of (ω, ψ_-) on $\mathfrak{h}$, and by definition $\psi_- := -\iota_X\phi$, $\sigma := \phi - \psi_- \wedge \theta$, $\sigma := \omega \wedge \omega$ for $\psi_- \in \Lambda^3\mathfrak{h}^*$, $\omega \in \Lambda^2\mathfrak{h}^*$,then The pair (ω, ψ_-) has a nearly half-flat SU(3)-structures which $d\psi_- = k\omega^2$ for $k \in \mathbb{R} - \{0\}$ with the torsion type $W_1^{\pm} \oplus W_2^- \oplus W_3$.*

Proof.

(I) Contraction of ψ_- and $\sigma := \omega \wedge \omega$

- $\iota_X\psi_- = \iota_X(-\iota_X\phi) = -\iota_X\iota_X\phi = 0$
- $\iota_X\sigma = \iota_X\phi - \iota_X(\psi_- \wedge \theta) = -\psi_- - (\iota_X\psi_- \wedge \theta - \psi_- \wedge \iota_X\theta) = -\psi_- + \psi_- \wedge \iota_X\theta = -\psi_- + \psi_- = 0$
- $X = \theta^\sharp$ *and* $\iota_X\theta = \theta(X) = 1$ *where* $\theta(Y) = g(X, Y)$ $\forall Y \in \mathfrak{g}$
  $\theta(X) = g(X, X) = |X|^2$

(II) The derivative of 3-form ψ_- and $\sigma := \omega \wedge \omega$ are as follow, and it admits nearly half-flat SU(3)-structures.

- $d\psi_- = d(-\iota_X\phi) = \iota_X(d\phi) = \iota_X(\theta \wedge \phi) = (\iota_X\theta) \wedge \phi - \theta \wedge (\iota_X\phi) = \theta(X)\phi - \theta \wedge (-\psi_-) = \phi + \theta \wedge \psi_- = \sigma + \psi_- \wedge \theta + \theta \wedge \psi_- = \sigma \Longrightarrow d\psi_- = \sigma = \frac{1}{2}\omega^2 \Longrightarrow k = \frac{1}{2}$, $k \in \mathbb{R} - \{0\}$
- $d\sigma = d(\phi - \psi_- \wedge \theta) = d\phi - d(\psi_- \wedge \theta) = d\phi - d\psi_- \wedge \theta + \psi_- \wedge d\theta = \theta \wedge \phi + \theta \wedge d\psi_- = \theta \wedge (\phi + d\psi_-) = \theta \wedge (\phi + \sigma)$

Theorem 1 . *Let $\mathfrak{h}$ be a six-dimensional nilpotent Lie algebra admitting a Nearly Half-Flat SU(3)-structures given by the pair (ω, ψ_-),and a closed Lee form, θ, such that $\theta(X) = 1$, $\forall X \in \mathfrak{z}(\mathfrak{h})$,then the seven-dimensional Lie algebra $\mathfrak{g} = \mathfrak{h} \oplus \mathbb{R}X$ admits a Locally Conformal Cocalibrated G_2-structures.*

Proof. *For a four-form $\phi = \frac{1}{2}\omega^2 + \psi_- \wedge \theta$, $\phi \in \Lambda^4\mathfrak{g}^*$ Look at parts (I) and (II) of preposition 1.*

Corollary 1. Those six-dimensional Lie algebras $\mathfrak{h}$ with double half-flat which are classified in [1] are admitting nearly half-flat SU(3)-structures for $X \in \mathfrak{z}(\mathfrak{h})$, then $\mathfrak{g} = \mathfrak{h} \oplus \mathbb{R}X$ where $\phi \in \Lambda^4\mathfrak{g}^*$ is Locally Conformally Cocalibrated G_2-structures, if $\mathfrak{g} = 1A, 1B, 1C, 2C, 2B$ and $3A$.

$$
\begin{aligned}
1A &= (0, 0, 0, e^{12}, e^{13}, e^{23}) \\
3A &= (0, 0, 0, 0, e^{12}, e^{15+34}) \\
1B &= (0, 0, e^{12}, e^{13}, e^{23}, e^{14}) \\
2B &= (0, 0, e^{12}, e^{23}, e^{14+35}) \\
1C &= (0, 0, e^{12}, e^{13}, e^{23}, e^{14+25}) \\
2C &= (0, 0, e^{12}, e^{13}, e^{23}, e^{14-25})
\end{aligned}
$$

Proof. The set of double half-flat six-dimensional lie algebras are subset of nearly half-flat six-dimensional lie algebras as any lie algebra $\mathfrak{h}_o$ with nearly half-flat SU(3)-structures has a equality in which $d\psi_- = \frac{1}{2}(\omega \wedge \omega)$, where in the case of double half-flat we have this equality in pulse : $d\psi_+ = 0$.
(section 4 of paper [1])

This allows ones to conclude that there exist L.C.CC G_2-structures on seven dimensional decomposable nilpotent lie algebras of the form $\mathfrak{g} = \mathfrak{h} + \mathbb{R}e_7$ where $\mathfrak{h}$ admits double half-flat lie algebras, so we are able to state the existence result for six double half-flat SU(3)-structures.But these are first result for building L.C.CC G_2-structures, as up to now we didn't find a stable 4-form ϕ_0 compatible with the orientation and metric g and it could be the next step on this paper.

3 Associated Lie algebras

Giving a method to obtain new 7-dimensional lie algebras endowed with locally conformally co-calibrated (L.C.CC) G_2-structures. Starting from 6-dimensions lie algebras with the nearly half-flat SU(3)-structures and then describing some explicit examples of them.
A SU(3)-structure on a lie algebra $\mathfrak{h}$ of dimension six, consist in a triple (g, J, Ψ) that (g, J) are almost Hermitian structure on $\mathfrak{h}$ and $\Psi = \psi_+ + i\psi_-$ is a complex volume $(0,3)$-form and satisfying :

$$\frac{1}{6!}\omega^6 = (-1)^3(\frac{i}{2})^3\psi \wedge \bar{\psi}$$

The *Käler* form associated to (ω, ψ_-) such that describe a metric as

$$g(X,Y)\omega^3 = -3\iota_X\omega \wedge \iota_Y\psi_- \wedge \psi_-$$

with $X, Y \in \mathfrak{h}$ and ι_X denoting the contraction by X, when (g, J, ψ_-) is a nearly half-flat SU(3)-structures on lie algebras $\mathfrak{h}$, we may choose an orthonormal frame $\{e_1, e_2, ..., e_6\}$ for almost complex structure s.t. $\psi_- = J\psi_+$ with the $\{e^1, e^2, ..., e^6\}$ an orthonormal dual basis and form ω and 3-form ψ_- can be written as $d\omega = 0$, $\Psi = \psi_+ + i\psi_-$

Now consider ϕ is a locally conformally co-calibrated G_2-structures and restricting the condition to decomposable cases s.t.

$$\mathfrak{g} = \mathfrak{h} \oplus \mathbb{R}$$

again consider $\{e_1, e_2, ..., e_6\}$ the basis for six-dimension vector space V and $\{e^1, e^2, ..., e^6\}$ basis for dual space V^* ,with one form $\theta \in \Lambda^1 V^*$ and pair of (ω, ψ_-) nearly half-flat SU(3)-structures where $\omega \in \Lambda^2 V^*$ and $\psi_- \in \Lambda^3 V^*$ on six-dimension lie algebras $\mathfrak{h}$, for 20 possible different ways to have 3-dimension basis as $\frac{6!}{3!3!} = 20$

$$\begin{array}{ccccc} e^{123} & e^{124} & e^{125} & e^{126} & e^{134} \\ e^{135} & e^{136} & e^{145} & e^{146} & e^{156} \\ e^{234} & e^{235} & e^{236} & e^{245} & e^{246} \\ e^{256} & e^{345} & e^{346} & e^{356} & e^{456} \end{array}$$

Where $e^{ijk} = e^i \wedge e^j \wedge e^k$ for $i, j, k \in \{1, 2, ..., 6\}$, are ordered as $i < j < k$

$$\psi_- := \sum_{i<j<k} \psi_{ijk} e^{ijk}$$

.

Is a generic 3-form. For four-form $\varphi \in \Lambda^4 V^*$ where V^* is dual space of V in seven-dimension with the basis $\{e^1, e^2, ..., e^6, , e^7\}$ and $\{e_1, e_2, ..., e_6, e_7\}$ respectively, so all of the possibilities for different basis would be $\frac{7!}{3!4!} = 35$

$$\begin{array}{ccccc} e^{1234} & e^{1235} & e^{1236} & e^{1237} & e^{1245} \\ e^{1246} & e^{1247} & e^{1256} & e^{1257} & e^{1267} \\ e^{1345} & e^{1346} & e^{1347} & e^{1356} & e^{1357} \\ e^{1367} & e^{1456} & e^{1457} & e^{1467} & e^{1567} \\ e^{2345} & e^{2346} & e^{2347} & e^{2356} & e^{2357} \\ e^{2367} & e^{2456} & e^{2457} & e^{2467} & e^{2567} \\ e^{3456} & e^{3457} & e^{3467} & e^{3567} & e^{4567} \end{array}$$

Two examples of L.C.CC G_2-structures:

First consider the lie algebra $1A = (0, 0, 0, e^{12}, e^{13}, e^{23})$ and one form

$$\theta := a_1 e_1 + a_2 e_2 + a_3 e_3 + a_4 e_4 + a_5 e_5 + a_6 e_6 + a_7 e_7$$

where $\theta \in \Lambda^1 V^*$ then the 4-form Φ when the generic form is :

$$\phi := \sum_{i<j<k<l} \phi_{ijkl} e^{ijkl}$$

and $\phi \in \Lambda^4 V^*$

$\phi = \phi_{3567}e^{3567} + \phi_{1467}e^{1467} + \phi_{1567}e^{1567} + \phi_{2345} + \phi_{2347}e^{2347} + \phi_{2357}e^{2357} + \phi_{2367}e^{2367} + \phi_{2467}e^{2467} + \phi_{2567}e^{2567} + \phi_{1257}e^{1257} + \phi_{1267} + \phi_{1347}e^{1347} + \phi_{1357}e^{1357} + \phi_{1367}e^{1367} + \phi_{1457}e^{1457} + \phi_{1234}e^{1234} + \phi_{1235}e^{1235} + \phi_{1236}e^{1236} + \phi_{1237}e^{1247} + a_7\phi_{2345}e^{4567} + (-a_7\phi_{1235} + \phi_{1567})e^{3457} + (-a_7\phi_{1236} + \phi_{2567})e^{3467} + (-a_7\phi_{1234} + \phi_{1467})e^{2457} + \phi_{2345}e^{1256} - \phi_{2345}e^{1346}$

This is a 4-form in seven dimension with the real lie algebra 1A, this form accepted locally conformal cocalibrated (L.C.CC) G_2-structure.

Now consider third , $1B = (0, 0, e^{12}, e^{13}, e^{14})$ and one form θ as previous again:

$$\theta := a_1 e_1 + a_2 e_2 + a_3 e_3 + a_4 e_4 + a_5 e_5 + a_6 e_6 + a_7 e_7$$

$\phi = (a_7\phi_{1236} + \phi_{1567})e^{2467} + \phi_{2567}e^{2567} + (a_7\phi_{1245} + \phi_{2567})^{3457} + (a_7\phi_{1246})e^{3467} + \phi_{1457}e^{1457} + \phi_{1467}e^{1467} + \phi_{1567}e^{1567}\phi_{2567}e^{2567} + \phi_{1257}e^{1257} + \phi_{1267} + \phi_{1347}e^{1347} + \phi_{1357}e^{1357} + \phi_{1367}e^{1367} + \phi_{1457}e^{1457} + \phi_{1234}e^{1234} + \phi_{1235}e^{1235} + \phi_{1236}e^{1236} + \phi_{1237}e^{1247} + a_7\phi_{2345}e^{4567} + (-a_7\phi_{1235} + \phi_{1567})e^{3457} + (-a_7\phi_{1236} + \phi_{2567})e^{3467}$

Acknowlegment: I would like to thanks Alberto Raffero and Leonardo Bagaglini and very special thanks to Anna Maria Fino for giving the problem and her suggestions, also I am thankful for Sigbjørn Hervik for his interest on my work.

Contact. Dipartamento di matematica,Via Carlo Alberto 10,
Università degli studenti di Torino, Italy. E-mail: azohrabi@unito.it

References

[1] Simon Chiossi,Andrew Swann, *G_2-structures with torsion form half-integrable nilmanifold*, arXiV:math/0404554

[2] Marisa Fernández, Anna Fino, Alberto Raffero, *Locally conformal calibrated G2-manifolds*, arXiv:math/1504.04508

[3] Leonardo Bagaglini , *Invariant cocalibrated G_2-structures on nilmanifolds* , arXiv:math/1509.01971

[4] A.Gray, M.Harvella , *the sixteen classes of almost hermitian manifolds and their linear invariant* (1978)

[5] Anna Fino, Alberto Raffero, *Einstein locally conformal calibrated G_2-structures*, arXiV:math/1303.6137

[6] Anna Fino, Alberto Raffero, *Coupled SU(3)-structures and Supersymmetry*, arXiV:math/1505.01655

[7] Diego Conti,Marisa Fernández, *Nilmanifolds with a calibrated G_2-structure* , arXiV:math/1008.0797

[8] Diego Conti, *Half-flat nilmanifolds*, arXiV:math/0903.1175

[9] Robert Bryant, *Remarks on the geometry of almost complex 6-manifolds*, arXiv:math/0508428

[10] Robert Bryant, *Some remarks on G_2-structures*, arXiv:math/0305124

[11] Ilka Agricola, Simon Chiossi,Anna Fino, *Solvmanifolds with integrable and non-integrable G_2-structures*, arXiv:math/0510300